

Emory University Hospital

Stroke Performance Measures

Quality Metric	Description	Bench-mark	2023	2024
VTE Prophylaxis	This measure captures the proportion of ischemic or hemorrhagic stroke patients who received VTE prophylaxis or have documentation why no VTE prophylaxis was given the day of or the day after hospital admission	85%	98%	98%
D/C on Antithrombotic Therapy	Percent of patients with an ischemic stroke or TIA prescribed antithrombotic therapy at discharge.	85%	100%	99%
Anticoagulation for Afib	Percent of patients with an ischemic stroke or TIA with atrial fibrillation/flutter discharged on anticoagulation therapy.	85%	100%	99%
Arrive by 2 Treat by 3	Percent of acute ischemic stroke patients who arrive at hospital within 2 hours of time last known well and for whom IV t-PA was initiated within 3 hours of time last known well.	85%	100%	100%
Early Antithrombotic	Percent of patients with ischemic stroke or TIA who receive antithrombotic therapy by the end of hospital day two.	85%	97%	97%
Discharged on Statin	Percentage of Ischemic Stroke and TIA patients who are prescribed high-intensity statin therapy at discharge OR, if >75 years of age, are prescribed at least moderate-intensity statin therapy at discharge.	85%	100%	99%
Stroke Education	Percent of patients with Stroke or TIA or their caregivers who were given education and/or educational materials during the hospital stay addressing ALL of the following: personal risk factors for stroke, warning signs for stroke, activation of emergency medical system, need for follow-up after discharge, and medications prescribed.	85%	99%	97%
Rehab Considered	Percent of patients with Stroke who were assessed for rehabilitation services.	85%	99%	99%
NIHSS for Ischemic Stroke	Percent of ischemic stroke and stroke not otherwise specified patients with a score reported for NIH Stroke Scale (Initial)	85%	98%	97%
Symptomatic Carotid Stent 30-Day Stroke or Death	30-day stroke or death rate following a percutaneous carotid stent procedure for symptomatic carotid stenosis	≤6%	0%	0%
Symptomatic Carotid Endarterectomy 30-Day Stroke or Death	30-day stroke or death rate following a surgical carotid endarterectomy procedure for symptomatic carotid stenosis	≤6%	0%	0%
Symptomatic TransCarotid Artery Revascularization 30-Day Stroke or Death	30-day stroke or death rate following a surgical carotid stenting procedure for symptomatic carotid stenosis	≤6%	0%	0%